

DFPD/98/TH/03

hep-th/9801038

Revised v2

NEW SUPERSYMMETRIC VACUA FOR D=4, N=4 GAUGED SUPERGRAVITY ¹

HARVENDRA SINGH ²

INFN Sezione di Padova, Dipartimento di Fisica ‘Galileo Galilei’,
Via F. Marzolo 8, 35131 Padova, Italy

ABSTRACT

In this paper we obtain supersymmetric brane-like configurations in the vacuum of $N = 4$ gauged $SU(2) \times SU(2)$ supergravity theory in four space-time dimensions. Almost all of these vacuum solutions preserve either half or one quarter of the supersymmetry in the theory. We also study these solutions in presence of nontrivial axionic and gauge field backgrounds. In the case of pure gravity with axionic charge the geometry of the spacetime is $AdS_3 \times R^1$ with $N = 1$ supersymmetry. An interesting observation is that the domain walls of this theory cannot be given an interpretation of a 2-brane in four dimensions. But it still exists as a stable vacuum. This feature is quite distinct from the domain-wall configuration in massive type IIA supergravity in ten dimensions.

¹ Research supported by INFN fellowship

²e-mail: hsingh@pd.infn.it

1 Introduction

Recent understanding of the nonperturbative aspects in string theories and its duality symmetries [1] has triggered a series of new developments in the study of massive supergravity theories in various dimensions [2, 3, 4, 5, 6, 7, 8, 9, 10]. It is viewed that the massive supergravities are required for the spacetime interpretations of higher dimensional world volume Dirichlet objects (D-p-branes) of type II superstrings. For example, massive (gauged) type IIA supergravity [2] is a candidate theory for type IIA D-8-brane [11, 3, 12] whose field strength is a 10-form. The dual of a 10-form in ten dimensions is a constant which can be identified as a mass term (cosmological constant) of massive type IIA supergravity. Next, the strong coupling limit of type IIA strings is M-theory whose spacetime theory is sought to be the eleven dimensional supergravity [13]. String dualities require that all even branes of type IIA and odd branes of IIB should have an eleven-dimensional interpretation [14]. In this picture a D-8-brane can be obtained from a double dimensional reduction of M-9-brane or toroidal reduction of an M-8-brane. Therefore 11-dimensional supergravity should also have massive analog of type IIA supergravity. There has been some progress recently in finding massive 11-dimensional supergravity [9, 10].

There are various ways to get massive supergravities starting from the massless ones. For example the generalised Scherk and Schwarz dimensional reduction scheme [15] do give rise to massive supergravities. Under generalised reduction scheme some of the higher dimensional fields are given linear dependences along the directions to be compactified [3, 4]. This procedure has been used in [3] to demonstrate the type IIA and IIB duality in $D=9$ for the massive case. Other methods require the gauging of internal global symmetries in the supergravity theory [16, 17]. An interesting feature of massive or gauged supergravity theories is that, in general, they possess original amount of supersymmetry even in the presence of mass (gauge) parameters. If supersymmetry is only the criterion then massive supergravities are as consistent as massless ones. However, distinction appears in terms of internal symmetries. The price we pay is that the internal global symmetries including dualities are generally reduced to smaller symmetry groups during the process of gauging. Once the mass (or gauge) parameters are set to zero the ‘standard’ (massless) supergravities are recovered.

In this work we will study $N = 4$ gauged $SU(2) \times SU(2)$ supergravity in four dimensions [17]. It has been recently established that this gauged supergravity can be

embedded into $N = 1$ supergravity in ten dimensions as an $S^3 \times S^3$ compactification [22]³. Previously also, a Kaluza-Klein(KK) interpretation for the $SU(2) \times SU(2)$ gauged supergravity was given in [23] where this model was identified as part of the effective $D = 4$ field theory for the heterotic string theory in an $S^3 \times S^3$ vacuum. These two KK interpretations are essentially the same upto consistent truncations. Thus it is worthwhile to study above $N = 4, D = 4$ gauged supergravity keeping in mind the recent developments in massive supergravities.

It is well known that the dimensional reduction of ten-dimensional supersymmetric Yang-Mills theory to 4 dimensions gives us $SU(4)$ invariant supersymmetric theory[18]. An off-shell supergravity with $SU(4)$ global symmetry and $N = 4$ local supersymmetry was constructed in four dimensions long ago [19]. The spectrum of this theory in the gravity multiplet consists; the graviton E_μ^M , 4 Majorana spin-3/2 gravitinos Ψ_μ^I , 3 vector fields A_μ^a , 3 axial-vector fields B_μ^a , 4 spin-1/2 Majorana fields χ^I , the dilaton field ϕ , and a pseudo-scalar field η . There do exist other versions of $N = 4$ supergravity in $D = 4$, e.g. $SO(4)$ supergravity theory[20]. It turns out that at the equation of motion level both the above supergravities are equivalent upto certain field redefinitions and duality relations. The internal subgroup $SU(2) \times SU(2)$ of $SU(4)$ supergravity can be gauged, giving rise to $N = 4$ supergravity with local $SU(2) \times SU(2)$ symmetry [17]. However, imposition of local supersymmetry in presence of the local internal symmetry requires introduction of mass like terms (or cosmological constant) in the theory in addition to certain bilinear fermionic terms [17]. The mass term appears as an effective dilaton potential in the theory. It was noted [17] that the dilaton potential leads to energy density which is unbounded from below which is physically nonacceptable⁴. But the question can be asked whether this theory yields some stable vacuum configurations. It has been noted that theory allows supersymmetric solutions like ‘electro-vac’ solutions[21] and more recently non-Abelian solitons as stable vacuum configurations were also shown to exist[24, 22]. In related work on strings in curved backgrounds [25], exact supersymmetric solutions of four-dimensional gauged supergravities have been constructed using the powerful techniques of (super)conformal field theory by exploiting a connection between gauged supergravities and non-critical strings.

In this work we will show that extended objects like strings, domain walls, pure axionic gravity, dilaton-axion gravity and maximally symmetric point-like configura-

³I thank Prof. P.K. Townsend bringing this fact into my knowledge

⁴ Similar unbounded potential also arises in gauged $SO(4)$ supergravity theory [16]

tions do exist in the vacuum of the $N = 4, D = 4$ gauged theory. All these backgrounds preserve half or one-fourth of the supersymmetries except for the point-like configuration for which we have not been able to find supersymmetric case. We also present the generalisation of the fundamental string solution as a vacuum solution of the massive supergravity. The paper has been organised as follows. In the next section we briefly describe the gauged model. In the subsequent sections III, IV, and V we obtain various background solutions and discuss there supersymmetric properties in each case. In the last section VI we summarise our results.

2 The Gauged Model

We consider the truncated version of the $N = 4$ gauged $SU(2)_A \times SU(2)_B$ supergravity model [17] given by the condition $B_\mu^a = 0$, that is half of the gauge fields are vanishing. In addition we shall only study the configurations which are either purely electric type or magnetic type with respect to remaining 2-form field strength. We also set all the spinor fields to zero in our study. Under these specifications the action [17] becomes

$$S = \int d^4x \sqrt{-g} \left[R + \frac{1}{2} \left(\partial_\mu \phi \partial^\mu \phi + e^{2\phi} \partial_\mu \eta \partial^\mu \eta \right) - \frac{1}{2 \cdot 2!} e^{-\phi} F_{\mu\nu}^a F^{a\mu\nu} - \frac{1}{2 \cdot 2!} \eta F_{\mu\nu}^a \tilde{F}^{a\mu\nu} + \frac{\Lambda^2}{2} e^\phi \right] \quad (1)$$

where $\Lambda^2 = e_A^2 + e_B^2$, e_A and e_B being the gauge couplings of respective $SU(2)$ groups and

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + e_A \epsilon_{abc} A_\mu^b A_\nu^c, \\ \tilde{F}_{\mu\nu}^a = \frac{1}{2} \epsilon_{\mu\nu}^{\sigma\rho} F_{\sigma\rho}^a.$$

The corresponding supersymmetry transformations are [17],

$$\delta \bar{\chi}^I = \frac{i}{2\sqrt{2}} \bar{\epsilon}^I \left(\partial_\mu \phi + i\gamma_5 e^\phi \partial_\mu \eta \right) \gamma^\mu - \frac{1}{4} e^{-\frac{\phi}{2}} \bar{\epsilon}^I \alpha^a F_{\mu\nu}^a \Gamma^{\mu\nu} + \frac{1}{4} e^{\frac{\phi}{2}} \bar{\epsilon}^I (e_A + i\gamma_5 e_B), \\ \delta \bar{\Psi}_\mu^I = \bar{\epsilon}^I \left(\bar{\partial}_\mu - \frac{1}{2} \omega_{\mu,mn} \Gamma^{mn} + \frac{1}{2} e_A \alpha^a A_\mu^a \right) - \frac{i}{4} e^\phi \bar{\epsilon} \gamma_5 \partial_\mu \eta \\ - \frac{i}{4\sqrt{2}} e^{-\frac{\phi}{2}} \bar{\epsilon}^I \alpha^a F_{\nu\rho}^a \gamma_\mu \Gamma^{\nu\rho} + \frac{i}{4\sqrt{2}} e^{\frac{\phi}{2}} \bar{\epsilon}^I (e_A + i\gamma_5 e_B) \gamma_\mu, \\ \delta(bosons) = 0, \quad (2)$$

where ϵ^I are four spacetime dependent Majorana spinors. Since fermionic backgrounds are absent therefore the bosonic fields do not vary under supersymmetry.

Our convention for the metric is with mostly minus signs $(+ - - -)$, $\omega_\mu{}^{mn}$ are the spin connections, $\{\gamma_\mu, \gamma_\nu\} = 2 g_{\mu\nu}$, $(\mu = 0, 1, 2, 3)$, and $\Gamma_{mn} = \frac{1}{4}[\Gamma_m, \Gamma_n]$, where m, n are tangent space indices. α^a are three 4×4 $SU(2)_A$ matrices and along with other three matrices of $SU(2)_B$ generate the $(\frac{1}{2}, \frac{1}{2})$ representation of the gauged model[17].

3 Domain Walls/Membranes

3.1 Pure dilaton gravity

The first solution we obtain is a domain wall in four dimensions with the following configuration satisfying the field equations derived from the action (1),

$$\begin{aligned} ds^2 &= U(y) \left(dt^2 - dx_1^2 - dx_2^2 \right) - U(y)^{-1} dy^2, \\ \phi &= -\ln U, \quad U = m|y - y_0|, \\ A_\mu^a &= 0, \quad \eta = 0, \end{aligned} \tag{3}$$

where $m^2 = \Lambda^2/2$. This background is singular at $y = y_0$ which is the position of the $t - x$ hyperplane or domain wall. Since no matter fields are present other than the dilaton this background represents pure dilaton gravity. Note that e^ϕ , analog of string coupling, vanishes at asymptotic infinity ($y \rightarrow \pm\infty$) and so also the curvature scalar. But both are divergent at $y = y_0$. The isometry group for this background is $P_3(1, 2) \times \mathbb{Z}_2$, where P_3 represents three dimensional Pöncare group and $\mathbb{Z}_2$ is the reflection symmetry of the dilaton potential around $y = y_0$.

We now study the supersymmetric properties of this background. If we substitute (3) in the supersymmetry equations (2) we find that the fermionic variations vanish provided the supersymmetry parameters satisfy

(a) when $e_B = 0$, $\Lambda = e_A$

$$\begin{aligned} \bar{\epsilon}^I \Gamma_3 &= i \bar{\epsilon}^I, \quad \text{for } U = m|y - y_0|, \\ \bar{\epsilon}^I &= U^{\frac{1}{4}} \bar{\epsilon}_0^I \end{aligned} \tag{4}$$

(b) and when $e_A = 0$, $\Lambda = e_B$

$$\begin{aligned} \bar{\epsilon}^I \Gamma_3 \gamma_5 &= - \bar{\epsilon}^I, \\ \bar{\epsilon}^I &= U^{\frac{1}{4}} \bar{\epsilon}_0^I \end{aligned} \tag{5}$$

where ϵ_0^I is a constant spinor. The conditions (4) and (5) can be satisfied because for spacial indices all Γ 's have imaginary eigen values $\pm i$, while $\gamma_5^2 = 1$. These

conditions break half of the supersymmetries in either case. Thus we see that there exist nontrivial Killing spinors preserving $N = 2$ supersymmetry for pure dilatonic domain wall backgrounds which is sufficient for the stability of the solution against quantum fluctuations.

3.2 Pure axionic gravity

A nontrivial axionic field configuration can also be obtained in the vacuum of this theory. Let us first present a much familiar geometry which is the solution of the field equations derived from (1),

$$\begin{aligned} ds^2 &= \left((1 + Qr^2)dt^2 - \frac{dr^2}{(1 + Qr^2)} - r^2 d\theta^2 \right) - dy^2, \\ \phi &= 0, \quad A_\mu^a = 0, \\ \eta &= \pm 2\sqrt{Q} y, \end{aligned} \tag{6}$$

such that $4Q = \Lambda^2/2$. The geometry of the four manifold is an anti-de Sitter (AdS) space $AdS_3 \times R$ ⁵. This solution is analogous to the Freedman-Gibbons (FG) pure electro-vac solution [21] which has the geometry of $AdS_2 \times R^2$, we will discuss it in the next section. We expect this background to be stable if the stability analysis is done based on the methods of anti-de Sitter spaces [26]. Instead we try to show here that the solution (6) will be stable from the supersymmetry arguments. We present another solution below which has similar asymptotic properties as the background in (6). We write down a new axionic gravity vacuum solution of (1)

$$\begin{aligned} ds^2 &= \left(f^2 dt^2 - \frac{dr^2}{f^2} - f^2 d\theta^2 \right) - dy^2, \\ \phi &= 0, \quad f(r) = 1 + \sqrt{Q} r \\ A_\mu^a &= 0, \quad \eta = 1 \pm 2\sqrt{Q} y, \end{aligned} \tag{7}$$

with $4Q = \Lambda^2/2$ where $2\sqrt{Q}$ is the measure of the axion charge. Note that for $Q = 1$ the asymptotic properties of both the solutions (6) and (7) are the same. But the background in (7) is much transparent from the supersymmetry point of view. If we substitute (7) in eq.(2) we find that the background preserves 1/4 of the

⁵ When y-coordinate is a closed circle the geometry will be $AdS_3 \times S^1$.

supersymmetries. Corresponding Killing spinors are for $e_A = 0$

$$\begin{aligned}\epsilon^I &= f(r)^{\frac{1}{2}} \epsilon_0^I \\ \bar{\epsilon}^I \Gamma_3 &= \pm i \bar{\epsilon}^I, \quad \bar{\epsilon}^I \Gamma_1 \gamma_5 = -\bar{\epsilon}^I.\end{aligned}\tag{8}$$

The twin conditions in (8) on the Killing spinors break the supersymmetry to 1/4th. Thus pure axionic gravity in the gauged model has $N = 1$ supersymmetry intact. Other conditions when $e_B = 0$, $\Lambda = e_A$ can also be similarly derived.

3.3 Nonvanishing dilaton and axion

Now we generalise above two special cases and show that both dilaton and axion can exist together in the vacuum of the gauged supergravity in a supersymmetric fashion. We obtain correspondingly a solution

$$\begin{aligned}ds^2 &= U(y) \left(f^2 dt^2 - \frac{dr^2}{f^2} - f^2 d\theta^2 \right) - U(y)^{-1} dy^2, \\ \phi &= -\ln U, \quad U(y) = 1 + m|y - y_0|, \\ \eta &= 1 + 2\sqrt{Q} y, \quad f(r) = 1 + \sqrt{Q} r \\ A_\mu^a &= 0,\end{aligned}\tag{9}$$

such that $m^2 = -4Q + \Lambda^2/2$. This solution is regular at $y = y_0$ as compared to domain wall solution in (3)⁶. The geometry of the manifold is $AdS_3 \times R$. Thus the Minkowskian flat geometry in (3) has turned into an anti-deSitter geometry in presence of nontrivial axion flux. The corresponding isometry group now is $O(2, 2)$ of the AdS_3 , Z_2 symmetry is broken due to nontrivial axion field. This background preserves 1/4-th of the supersymmetries. We find that most general Killing spinors can be written when $m^2 = e_A^2/2$ and $4Q = e_B^2/2$,

$$\begin{aligned}\epsilon^I &= U(y)^{\frac{1}{4}} f(r)^{\frac{1}{2}} \epsilon_0^I \\ \bar{\epsilon}^I \Gamma_3 &= i \bar{\epsilon}^I, \quad \bar{\epsilon}^I \Gamma_1 \gamma_5 = -\bar{\epsilon}^I.\end{aligned}\tag{10}$$

It is clear from (9) that pure dilatonic and pure axionic solutions are special cases of (9) when $Q = 0$ and $m = 0$ respectively. Note that in (9) only positive values for the axionic charges $\sqrt{Q}$ are allowed. This restriction comes from the supersymmetry in

⁶We could have as well chosen $U = 1 + m|y - y_0|$ in eq.(3) instead of taking a divergent solution at $y = y_0$.

presence of a regular dilaton field which is nontrivial now. While in the previous case of pure axionic gravity (no dilaton) both plus and minus values for the axion charge were allowed.

4 Vortices/Strings

4.1 Pure dilatonic Case

We also obtain vortex or string-like solution of the equations of motion derived from action (1). We look for the solutions such that the isometry group is $P_2(1, 1) \times SO(2)$. Again we consider pure dilaton gravity. The background solution is

$$\begin{aligned} ds^2 &= U(y)^2 (dt^2 - dx^2) - dy^2 - y^2 d\theta^2, \\ \phi &= -2 \ln U, \quad U = m y, \\ A_\mu^a &= 0, \quad \eta = 0, \end{aligned} \tag{11}$$

where the parameter m satisfies $m^2 = \Lambda^2/8$. Here $y = \sqrt{y_1^2 + y_2^2}$ is the radial distance in the transverse 2-dimensional Euclidean plane and θ is the azimuthal angle. It should be noted that the function $U(y)$ in (11) is of very specific nature which vanishes at $y = 0$. It indicates that the volume of the $t - x$ space shrinks to zero at $y = 0$ much like the size of the 2-sphere shrinking to zero at the centre of the Cartesian three-space. For this background the curvature scalar is singular at the origin $y = 0$ and also the string coupling while at asymptotic infinity both are vanishing.

Background in (11) preserves half of the supersymmetries of the theory. There exist a nontrivial Killing spinor for $e_B = 0$

$$\begin{aligned} \bar{\epsilon}^I \Gamma_2 &= i \bar{\epsilon}^I, \\ \epsilon^I &= U^{\frac{1}{2}} \epsilon_0^I, \quad \text{for } U = m y. \end{aligned} \tag{12}$$

where Γ_2 corresponds to the direction tangent to y . Note that these Killing spinors vanish at $y = 0$. Other Killing spinors when $e_A = 0$ can also similarly be derived.

4.2 Presence of nonvanishing electric field

As in the case of Domain-walls, we find here that a nontrivial electric flux can be switched on in the supersymmetric string solution (11). The electric field is given by

a constant electric field tangent to the x-direction. Before showing that we assume that the singularity at the origin is modified in presence of electric field. We consider that at the origin $U|_{y=0} = 1$ so that string coupling is unity at the origin. Advantage of this is that we can take the smooth limit $m \rightarrow 0$ in the background below. The limit corresponds to constant dilaton field. It will be clear soon. Under these specifications the background solution of action (1) in presence of electric field is

$$\begin{aligned} ds^2 &= U(y) \left(\left(1 + \frac{q^2}{2} x^2\right) dt^2 - \frac{dx^2}{\left(1 + \frac{q^2}{2} x^2\right)} \right) - dy_1^2 - dy_2^2, \\ \phi &= -\ln U, \quad U = 1 + m y^2, \\ A_\mu^a &= (q x, 0, 0, 0) \delta^{a3}, \quad \eta = 0, \end{aligned} \tag{13}$$

such that $m = \frac{\Lambda^2 - q^2}{8}$, where q is the measure of the electric field strength⁷. From (13) it is clear that we have chosen an specific direction in the isospin space thus breaking $SU(2)_A$ to $U(1)$. Note that now there is no singularity at $y = 0$ and asymptotically string coupling and curvature scalar both vanish. Though background (13) does not preserve any supersymmetry but we are free to consider the limit $m \rightarrow 0$, $q \rightarrow \Lambda$, $e_A = 0$. This is the advantage of choosing the potential $U = 1 + m r^2$. Now we can set $m = 0$ in (13). The vacuum solution in (13) then reduces to Freedman-Gibbons purely electric ‘electro-vac’ solutions. It is easy to follow this by defining the following relations

$$x = \frac{1}{q} \tan \rho, \quad t = \frac{\tau}{q}, \quad -\frac{\pi}{2} \leq \rho \leq \frac{\pi}{2}. \tag{14}$$

Eq. (13) becomes,

$$\begin{aligned} ds^2 &= \frac{1}{\Lambda^2 \cos^2 \rho} (d\tau^2 - d\rho^2) - dy_1^2 - dy_2^2, \\ \phi &= 0, \quad U = 1, \\ A_\mu^a &= (q x, 0, 0, 0) \delta^{a3}, \quad \eta = 0, \end{aligned} \tag{15}$$

which is the electro-vac solution of [21] corresponding to case $e_A = 0$, $B_\mu^a = 0$. This background has $N = 2$ supersymmetry [21] and can be described as the extremal (or supersymmetric) limit $m \rightarrow 0$ of the background in (13).

⁷ When we take $q \rightarrow 0$ the background (13) is still a vacuum solution

4.3 Presence of axion charges

In the presence of axion field we obtain a vacuum solution for the action (1) which is a generalisation of the fundamental string solution [27]. Such solutions (non-static) with an isometry group $O(1,1) \times SO(2)$ have already been obtained in a previous work [7] by the author. But here we are going to present the background solution which is static and has the symmetry $R^1 \times Z_2 \times SO(2)$. This background is

$$\begin{aligned} ds^2 &= \left(f(x) dt^2 - \frac{dx^2}{f(x)} \right) - U(y, x) (dy^2 + y^2 d\theta^2), \\ \phi &= -\ln U(y, x), \quad U = 1 - Q \ln |y - y_0| + m |x - x_0|, \\ f(x) &= 1 + m |x - x_0|, \quad \partial \eta = e^{-2\phi} H, \\ B_{01} &= U^{-1} \end{aligned} \tag{16}$$

where H represents antisymmetric field strength of $B_{\mu\nu}$ and $m^2 = \Lambda^2/2$. Note that this background has invariances under time translations, reflections about $x = x_0$ plane and rotations in the two dimensional y-plane. This background, however, does not preserve supersymmetries in (2). But it has two special limits; one when $Q = 0$ (16) reduces to the supersymmetric domain wall solution in (3), other when $m = 0$ background (16) becomes a fundamental string [27]. Thus we claim that the background (16) is the most simple generalisation of string geometry, though unstable, in presence of dilatonic potential (cosmological constant) as in (1). That is to say, background (16) represents a fundamental string whose vacuum is a domain-wall instead of a Minkowski space. The string vacuum (domain-wall) is stable but not the strings (geometric excitations which carry axionic charges) in this vacuum. The stable geometries in the domain-wall vacuum which carry axionic charges have already been presented in section III.

5 Point-like Solutions

To complete the analysis we also obtain maximally symmetric isotropic background solutions of (1). We have not been able to find a supersymmetric case for them. The solution we are going to present below is of very specific type that it allows only unit value of the magnetic charge. The solution is given by

$$ds^2 = f dt^2 - \frac{dr^2}{f} - U (d\theta^2 + \sin^2 \theta d\Phi^2),$$

$$\begin{aligned}\phi &= -\ln U, & U &= 1 + m r, & f(r) &= \frac{r}{r_0} \pm 1, \\ A_\mu^a &= (0, 0, 0, Q_m(1 - \cos \theta)) \delta^{a3}, & \eta &= 0\end{aligned}\tag{17}$$

with $m = \frac{\Lambda^2 + Q_m^2}{2} r_0$ such that $Q_m = 1$. Thus this solution exists only in the presence unit value of magnetic charge. For plus sign in $f(r)$ the solution is smooth every where and is asymptotically flat. But for minus sign there appears a nonsingular horizon at $r = r_0$. From the nonsingular horizon we mean by the finite size. However, this does not hide any singularity.

6 Summary

We summarise our results as follows:

In this work we have obtained a stable domain-wall solution in the purely dilatonic vacuum of gauged $N = 4$ supergravity. The solution preserves half of the supersymmetries. Then we obtained pure axionic gravity vacuum which preserves 1/4 of the supersymmetries of the theory. We also have obtained domain-walls in the presence of nontrivial axionic background. To our surprise these backgrounds also preserve one fourth of the supersymmetries and are therefore stable. As we find that the domain wall solution cannot be given an interpretation of a dual of a 4-form field strength in the action (1) because the dilatonic potential is unbounded from below. Though, in the case of type IIA gauged supergravity[2] the domain walls have an interpretation as a dual of a 10-form field strength of an 8-D-brane.

Next, we have obtained vortex (string-like) solutions which preserve half of the supersymmetries and therefore are stable solutions. Then we have generalised these solutions in presence of constant electric flux and showed that these solutions are generalisation of Freedman-Gibbons electro-vac background. Though these solutions do not preserve any supersymmetry but can be smoothly taken to their extremal (supersymmetric) limit which is an electro-vac background preserving $N = 2$ supersymmetries.

In last we obtained the maximally symmetric point-like solutions also. We have'nt found any fraction of supersymmetry being left unbroken for point case. It should be explored further if black holes could be found in gauged $SU(2) \times SU(2)$ supergravity.

In conclusion we have seen that the vacuum of the gauged supergravity is quite interesting and is rich with stable brane-like configurations.

Note Added: When this work was communicated I came to know from Prof. P. K. Townsend that another domain-wall solution has been obtained in [28]. We can see that this domain-wall is related to the domain-wall solution in (3) by going to a frame where the potential $U(y) \rightarrow 1/U(y)$. However, the string coupling diverges at asymptotic infinity in that case. One thing can also be seen from (3) that the dilaton potential $-\Lambda^2/2e^\phi$ is bounded because $e^\phi \rightarrow 0$ as $y \rightarrow \pm\infty$.

Acknowledgements: I would like to thank Mario Tonin for many useful discussions. I also thank anonymous referee for bringing up the previous works [23, 25].

References

- [1] C.M. Hull and P.K. Townsend, *Unity of Superstring Dualities*, **Nucl. Phys.** **B438** (1995) 109, [[hep-th/9410167](#)].
 E. Witten, **Nucl. Phys.** **B443** (1995) 85, [[hep-th/9503124](#)].
 J. Polchinski, *Dirichlet branes and Ramond-Ramond charges*, **Phys. Rev. Lett.** **75** (1995) 4724.
 A. Strominger and C. Vafa, **Phys. Lett.** **B379** (1996) 99, [[hep-th/9601029](#)].
 T.Banks, W. Fischler, S. Shenker and L. Susskind, *M-theory as A Matrix Model; A Conjecture*, **Phys. Rev.** **D55** (1997) 5112, [[hep-th/9610043](#)].
 For more details see the reviews and lectures;
 A. Sen, *Unification of String Dualities*, [[hep-th/9609176](#)].
 P.K. Townsend, *Four Lectures on M Theory*, [[hep-th/9612121](#)].
 J. Polchinski, *Lectures on D-branes* in ‘Fields, Strings, and Duality’ (TASI 96), eds. C. Efthimiou and B. Greene, World Scientific 1997, [[hep-th/9611050](#)].
 J. Schwarz, *Lectures on Superstrings and M-Theory Dualities* in ‘Fields, Strings, and Duality’ (TASI 96), eds. C. Efthimiou and B. Greene, World Scientific 1997, [[hep-th/9607201](#)].
 C. Vafa, *Lectures on Strings and Dualities*, [[hep-th/9702201](#)].
 J. Maldacena, *Black Holes in String Theory*, [[hep-th/9607235](#)]; *Black Holes and D-branes*, [[hep-th/9705078](#)]. T. Banks, *Matrix Theory*, [[hep-th/9710231](#)].
 D. Bigatti and L. Susskind, *Review of Matrix Theory*, preprint SU-ITP-97/51, [[hep-th/9712072](#)].
- [2] L.J. Romans, **Phys. Lett.** **B169** (1986) 374.
 J.L. Carr, S.J. Gates, and R.N. Oerter, **Phys. Lett.** **B189** (1987) 68.

- [3] E. Bergshoeff, M. de Roo, M. Green, G. Papadopoulos and P. Townsend, *Duality of type II 7-branes and 8-branes*, **Nucl. Phys.** B470 (1996) 113, [**hep-th**/9601150].
- [4] P. Cowdall, H. Lu, C. N. Pope, K.S. Stelle and P. K. Townsend, **Nucl. Phys.** B486 (1997) 49.
I. V. Lavrinneko, H. Lu and C. N. Pope, *From topology to generalised dimensional reduction*, CTP-TAMU-59/96, [**hep-th**/9611134],
- [5] J. Maharana and H. Singh, *On the compactification of type II string theory*, **Phys. Lett.** B408 (1997) 164, [**hep-th**/9505058].
- [6] E. Bergshoeff, M. de Roo and E. Eyras, *Gauged supergravity from dimensional reduction*, preprint UG-7/97, [**hep-th**/9707130].
- [7] H. Singh, *Macroscopic String-like Solutions in Massive Supergravity*, preprint DFPD/97/TH/48, Phys. Lett. B (in press), [**hep-th**/9710189].
- [8] E. Bergshoeff, P.M. Cowdall and P.K. Townsend, *Massive type IIA supergravity from the topologically massive D-2-brane*, preprint UG-6/97, [**hep-th**/9707139].
- [9] P.S. Howe, N.D. Lambert, and P.C. West, *A New Massive type IIA supergravity from Compactification*, [**hep-th**/9707139].
- [10] E. Bergshoeff, Y. Lozano, and T. Ortin, *Massive Branes*, preprint UG-8/97, [**hep-th**/9712115].
- [11] J. Polchinski and E. Witten, *Evidence for heterotic-type I String Duality*, **Nucl. Phys.** B460 (1996) 525, [**hep-th**/9510169].
- [12] E. Bergshoeff and M. B. Green, *The type IIA super-eight brane*, preprint VG-12/95,
M.B. Green, C.M. Hull and P.K. Townsend, [**hep-th**/9604119].
- [13] E. Cremmer, B. Julia, and J. Scherk, *Supergravity Theory in 11 dimensions*, **Phys. Lett.** B76 (1978) 409.
- [14] P.K. Townsend, *M-Theory from its Superalgebra* (Cargese Lectures 97), [**hep-th**/9712004].

- [15] J. Scherk and J. H. Schwarz, **Nucl. Phys.** B153 (1979) 61; **Phys. Lett.** B82 (1979) 60.
- [16] A. Das, M. Fischler and M. Rocek, ITP-SB-77-15, ITP-SB-77-38; **Phys. Rev.** D16 (1977) 3427.
- [17] D.Z. Freedman and J.H. Schwarz, **Nucl. Phys.** B137 (1978) 333.
- [18] F. Gliozzi, J. Scherk and D. Olive, **Nucl. Phys.** B122 (1977) 253.
L. Brink, J. Schwarz and J. Scherk, **Nucl. Phys.** B121 (1977) 253.
- [19] E. Cremmer, J. Scherk and S. Ferrara, *SU(4) invariant Supergravity Theory*, **Phys. Lett.** B74 (1978) 61.
- [20] A. Das, **Phys. Rev.** D15 (1977) 259.
E. Cremmer and J. Scherk, **Nucl. Phys.** B127 (1977) 253.
- [21] D.Z. Freedman and G.W. Gibbons, **Nucl. Phys.** B233 (1984) 24.
- [22] A.H. Chamseddine and M.S. Volkov, *Non-Abelian Solitons in $N = 4$ Gauged Supergravity and Leading Order String Theory*, preprint RU/97-5-B, [**hep-th**/9711181].
- [23] I. Antoniadis, C. Bachas and A. Sagnotti, **Phys. Lett.** B235 (1990) 255.
- [24] A.H. Chamseddine and M.S. Volkov, **Phys. Rev. Lett.** 79 (1997) 3343.
- [25] I. Antoniadis, S. Ferrara and C. Kounnas, **Nucl. Phys.** B421 (1994) 343.
- [26] P. Breitenlohner and D. Freedman, **Phys. Lett.** B115 (1982) 197; Ann. of Phys. 144 (1982) 249
G.W. Gibbons, C.M. Hull and P. Warner, **Nucl. Phys.** B218 (1983) 173
E. Witten, Comm. Math. Phys.80 (1981) 381.
- [27] A. Dabholkar, G. Gibbons, J. Harvey and F. Ruiz-Ruiz, **Nucl. Phys.** B340 (1990) 53.
- [28] P.M. Cowdall, *Supersymmetric Electrovacs in Gauged Supergravities*, preprint R/97-51, [**hep-th**/9710214].